J. Dispersion Science and Technology, 22(6), 609–611 (2001)

Author Index to Voume 22

A

Acharya, A., 551
Adair, J. H., 143
Ahmad, S., 261
Ahn, H.-J., 591
Aimaretti, M., 13
Al-Kashef, I., 111
Aserin, A., 269

B

Balasubramanian, R., 485
Baszkin, A., 313
Bell, N. S., 143
Benichou, A., 269
Bhattacharjee, C., 71
Bhattacharya, S. Ch., 323
Bjerregaard, S., 23
Blanzat, M., 167
Brancewicz, C., 491

C

Carlotti, M. E., 13, 381
Chaix, C., 473
Chen, D., 291
Chen, J., 203
Cheng, S., 291, 337, 535
Chibowski, St., 281
Chiraleu, F., 231
Cho, S.-B., 143
Cirkel, P. A., 211
Cordobés, F., 409
Csóka, I., 461

D

Das, B., 491
Das, K. P., 71
Dékány, G., 461
Delair, Th., 473
de Oliveira Tiera, V. A., 177
Donescu, A., 231
Donescu, D., 231
Dong, J., 57
Du, J., 337, 529

E

Eastoe, J., 597
Elaïssari, A., 473
El-Ashgar, N. M., 583
El-Astal, A., 431
El-Nahhal, I. M., 583
Erös, I., 461

F

Filipović-Vinceković, N., 221
Fontana, M., 211
Førdedal, H., 33
Franco, J. M., 409
Frokjaer, S., 23
Furukawa, H., 245
Fusulan, L., 231

G

Gallarate, M., 13, 381
Gallegos, C., 409

Ganachaud, F., 473
Gao, Y., 89, 197
Garti, N., 269
Ghosh, J. K., 323
Ghosh, K. K., 343
Graeffe, M., 491
Grillo, I., 597
Guan, D.-R., 349
Guo, R., 329, 443, 541
Guo, X., 329, 443

H

Harashima, A., 245
Hashizaki, K., 185, 191
He, X., 89, 197
Heenan, R. K., 597
Holmberg, K., 297, 569
Horvath, L., 221
Hou, Z., 255
Huang, L., 421
Huang, X., 563

I

Ishii, F., 97
Ismail, Z., 261

J

Jesionowski, T., 363
Jiang, B., 453, 529
Johnsen, E. E., 33
Jönsson, B., 387
Juan, D., 291

K

Kabalnov, A., 1
Kamyshny, A., 313
Kassim, A., 261
Katougi, Y., 185, 191
Kayali, I., 111, 431
Kim, B., 485
Kim, J.-C., 591
Kim, J.-H., 79
Kim, J.-M., 399
Koper, G. J. M., 211
Kumar, A., 245
Kunieda, H., 119, 245

L

Laayoun, A., 473
Labadie, L., 167
Lee, E.-J., 591
Lee, H.-J., 79
Lee, J.-Y., 79
Lee, K.-N., 79
Leijon, H., 569
Leubner, I. H., 125, 373
Li, G., 355, 563
Li, G.-Z., 349, 421
Li, Z., 255
Liao, G.-Z., 421
Liu, H., 203
Lou, A.-J., 349
Luo, L., 41

M

Magdassi, S., 313
Mihaljević, B., 221
Mohlin, K., 569
Moon, T. L., 167
Moros, J. E., 409
Moulik, S. P., 551
Mu, J.-H., 421

N

Nagasaka, Y., 97
Nan, Z., 103
Neumann, M. G., 177
Nyden, M., 297

O

Ogawa, N., 185, 191
Olsson, U., 119
Ozawa, K., 119

P

Park, S.-K., 591
Paszkiewicz, M., 281
Perez, E., 167
Perrotta, A. J., 143
Petcu, C., 231
Picatonotto, T., 381
Pichot, C., 473

Q

Qian, J., 541

R

Rang, M.-J., 591
Rasmussen, D. H., 491
Rico-Lattes, I., 167
Risović, D., 221
Rosenholm, J., 491

S

Sadtler, B., 485
Saito, Y., 185, 191
Sanyal, S. K., 551
Shin, J., 399
Söderberg, I., 23
Söderman, O., 387
Song, M.-E., 591
Steytler, D., 597
Ström, C., 387
Suhaimi, H., 261
Summers, M., 597

T

Taguchi, H., 185, 191
Texter, J., 499; (book reviews), 139, 311
Thompson, D. H., 399
Tiera, M. J., 177
Tiwary, L. K., 343
Tomaišić, V., 221
Toscano, A., 491
Trotta, M., 13
Tsouris, C., 57

U

Uddin, Md. H., 245
Urdahl, O., 33

V

Vasilescu, M., 231
Vermehren, C., 23
Vione, D., 381

W

Wang, H., 255
Wang, Y., 355
Wang, Z.-W., 349, 421
Wassenius, H., 297
Wei, A., 485

X

Xiang, Q., 103
Xiang, Y., 103, 453
Xie, J., 103, 337, 453, 529
Xu, G., 355, 563
Xu, H., 329
Xu, J., 485
Xu, K., 89, 197

Y

Yi, X.-Z., 349
Yu, J., 41, 203
Yuan, S., 355, 563

Z

Zeng, X., 103, 291, 337, 453, 529, 535
Zhang, L., 41, 563
Zhang, X., 443
Zhao, B., 203
Zhao, S., 41
Zheng, L.-Q., 421
Zou, A., 541
Zourab, S., 111, 431, 583

J. Dispersion Science and Technology, 22(6), 613–617 (2001)

Subject Index to Volume 22

A

Acetate groups, from polyvinyl alcohol, influence on adsorption and electrochemical properties of the TiO_2-polymer solution interface, 281–289
Acetylsalicylic acid, effect of microemulsion structures on the hydrolysis of, 541–549
Acidic components, effect on the dynamic interfacial tensions in surfactant/alkali/acidic crude oil systems, 41–55
Acrylic acid-co-ethyl methacrylate copolymers, fluorescence probe study of interaction with SDS, 177–184
Adsorption
  of a divalent cationic surfactant onto silica, influence of added salt on, 387–396
  Al^{3+} counterion, rheological properties of anionic surfactant solution in the presence of, 421–429
Alkyl ketene dimer, spontaneous emulsification of, 569–581
α-Alumina particles, morphological control when synthesized in 1,4-butanediol solution, 143–165
Aluminum hydroxide particles, nanoscale, precipitation of, 491–498
Amine-functionalized latex microspheres, grafting of oligodeoxyribonucleotides onto, 473–484
Anionic copolymer, structural and composition effect on protein flocculation, 203–209
Anthracene, fluorescence quenching by N,N-diethylaniline and phenothiazine in CTAB/n-$C_5H_{11}OH/H_2O$ microemulsion, 329–335
Applied electric fields, phase inversion of liquid–liquid dispersions under, 57–69

B

Betaine surfactant, effect of polyvinylpyrrolidone on the states of water in W/O microemulsions with, 563–567
Binuclear complexes, coordinating tripeptide in micellar solution for catalysis of the hydrolysis of *p*-nitrophenyl picolinate, 103–109
Biomolecules, hydrophobically modified, interfacial properties of, 313–322
1,4-Butanediol solution, morphological control of α-alumina particles synthesized in, 143–165

C

Calcium thioglycolate, formulation and characterization of W/O/W multiple emulsions of, 13–21
Carboxylic acid sugars, synthesis and physicochemical study of surfactants derived from, 167–176
Chemical cross linking, characterization of β-lactoglobulin microcapsules formed by, 71–78
Colloid science foundations (book review), 311–312
Complex coacervation, for preparation and characterization of triclosan-containing microcapsules, 591–596
Condensation, of organic particles, 499–528
Continuous stirred tank reactor crystallizer, seeded precipitations in, 373–380
Copper(II)-salicylate complex, effect of surfactant microenvironment on SOD-like activity of, 197–202
Cosurfactant, short chain, influence of monomers on microemulsions with, 231–244

Counterion specificity, effect on counterion binding to ionic micelles, 221–229
Cyclodextrins
solubilizing ability for (+)-α-pinene, 185–190
and surfactant mixed systems, solubilization of (+)-α-pinene by, 191–195

D

bis{N-(2-Deoxy-β-D-glucopyranosyl-2- [3-carboxyl salicylaldimino])} M2(II), for catalytic hydrolysis of *p*-nitrophenyl picolinate in CTAB micellar solution, 453–459
N,N-Diethylaniline, in CTAB/n- $C_5H_{11}OH/H_2O$ microemulsion, for fluorescence quenching of anthracene, 329–335
2,5-Dihydroxyl-1,4-benzoquinone, in aqueous solution, effects of surfactants on hydride transfer from methylene blue leudodye to, 535–539
Dispersions—Characterization, Testing, and Measurement (book review), 139–141
Dodecyl dimethyl amino oxide/SDS, phase behavior in the presence of perfluorooctanol, 111–117

E

Electrical double layer potential, for spherical particles, 349–354
Emulsification, spontaneous, of alkyl ketene dimer, 569–581
Emulsions
systems, preparation of silica particles in, 363–371
water-in-crude-oil under flowing conditions, measurement of viscosity in, 33–39
W/O
accelerated stability testing, 23–31
for food applications, polyols, high pressure, and refractive indices equalization for improved stability of, 269–280
W/O/W multiple, of calcium thioglycolate, 13–21
Encapsulation
efficiency, calculation for liposomes, 97–101
of phenolphthalein pH-indicator into a sol-gel matrix, 583–590
Experimental design, for evaluation of dispersion stability, 297–309

F

Flocculation, of protein, structural and composition effect of anionic copolymer on, 203–209
Fluorescence probe
2-methyl naphthalene, in ionic and nonionic micelles, 323–328
study, of the interaction between acrylic acid-co-ethyl methacrylate copolymers and SDS, 177–184
Fluorescence quenching
of anthracene by N,N-diethylaniline and phenothiazine in CTAB/n- $C_5H_{11}OH/H_2O$ microemulsion, 329–335
by halides, 323–328
Food applications, polyols, high pressure, and refractive indices equalization for improved stability of W/O emulsions for, 269–280

G

Gold nanoparticles, large, extraction and dispersion in nonpolar solvents, 485–489

H

Halides, fluorescence quenching by, 323–328
Heat setting, characterization of β-lactoglobulin microcapsules formed by, 71–78
Hydride transfer, from methylene blue leudodye to 2,5-dihydroxyl-1,4- benzoquinone in aqueous solution, effects of surfactants on, 535–539
Hydrolysis
of acetylsalicylic acid, effect of microemulsion structures on, 541–549
catalytic
of *bis*(*p*-nitrophenyl) phosphate by lipophilic diaquo(tetraamine) cobalt(III) in micellar solution, 337–341
of *p*-nitrophenyl picolinate by *bis*{N-(2- deoxy-β-D-glucopyranosyl-2-[3-carboxyl salicylal-dimino])} M_2(II) in CTAB micellar solution, 453–459
microemulsions as a reaction media for, 343–348
of *p*-nitrophenyl picolinate catalyzed by copper(II) and zinc(II) binuclear complexes coordinating tripeptide in micellar solution, 103–109
Hydrophobically modified biomolecules, interfacial properties of, 313–322

I

Interface electrophoresis method, for stability of polyacrylate dispersions, 291–295
Interfacial properties, of hydrophobically modified biomolecules, 313–322

Interfacial tensions
in surfactant/alkali/acidic crude oil systems, effect of acidic components on, 41–55
ultra-low, in oil-water-mixed surfactant systems, 255–259

K

Kaolinite, adsorption of sodium oleate on, 355–362

L

β-Lactoglobulin microcapsules, formed by chemical cross linking and heat setting, 71–78
Liposomes
estimation of marker entrapped in, 97–101
interaction with neutral polymers, 461–472
Liquid–liquid dispersions, phase inversion under applied electric fields, 57–69

M

Metal ions, effects on the micellization of ionic surfactants, 529–533
Metallomicellar catalysis
of the catalytic hydrolysis of *bis*(*p*- nitrophenyl) phosphate, 337–341
of the catalytic hydrolysis of *p*-nitrophenyl picolinate by *bis*{N-(2-deoxy-β-D-glucopyranosyl-2-[3-carboxyl salicylaldimino])} M_2(II) in CTAB micellar solution, 453–459
2-Methyl naphthalene, as a fluorescence probe in ionic and nonionic micelles, 323–328
Methylene blue leudodye, effects of surfactants on hydride transfer to 2,5-dihydroxyl-1,4-benzoquinone in aqueous solution, 535–539
Micelles
inverted cylindrical branched, structure and percolation of, 211–219
ionic, counterion binding to, 221–229
ionic and nonionic, 2-methyl naphthalene as fluorescence probe in, 323–328
Micellization, of ionic surfactants, effects of metal ions on, 529–533
Microcapsules
containing triclosan, preparation and characterization by complex coacervation, 591–596
β-lactoglobulin, formed by chemical cross linking and heat setting, 71–78
Microemulsions
biological system, using mixed oil (ricebran and isopropyl myristate), polyoxyethylene(2)oleyl ether (Brij 92), isopropyl alcohol, and water, 551–561
CTAB/n-$C_5H_{11}OH/H_2O$, fluorescence quenching of anthracene by N,N-diethylaniline and phenothiazine in, 329–335
nonionic, oil-induced structural changes in, 119–124
polymerization of cationic surfactant films in, 597–606
as reaction media for a hydrolysis reaction, 343–348
with short chain cosurfactant, influence of monomers on, 231–244
solubilization enhancing effect of A-B-type silicone surfactants in, 245–253
structures, effect on the hydrolysis of acetylsalicylic acid, 541–549
W/O, effect of polyvinylpyrrolidone on the states of water in, 563–567
Microenvironment, created by surfactant
effect on SOD activity, 89–96
effect on SOD-like activity of copper(II)-salicylate complex, 197–202
Microspheres
amine-functionalized latex, grafting of oligodeoxyribonucleotides onto, 473–484
phenolic/furfural aerogel, direct formation in supercritical carbon dioxide, 79–87
Mixed oil (ricebran and isopropyl myristate), polyoxyethylene(2)oleyl ether (Brij 92), isopropyl alcohol, and water biological microemulsion, 551–561
Monomers, influence on microemulsions with short chain cosurfactant, 231–244
Morphological control, of α-alumina particles synthesized in 1,4-butanediol solution, 143–165
Multicomponent systems, Ostwald ripening in, 1–12

N

Nanoparticles, large gold, extraction and dispersion in nonpolar solvents, 485–489
Neutral polymers, interaction with liposomes, 461–472
bis(*p*-Nitrophenyl) phosphate, catalytic hydrolysis by lipophilic diaquo(tetraamine) cobalt(III) in micellar solution, 337–341
p-Nitrophenyl picolinate hydrolysis
catalyzed by copper(II) and zinc(II) binuclear complexes coordinating tripeptide in micellar solution, 103–109

by *bis*{N-(2-deoxy-β-D-glucopyranosyl- 2-[3-carboxyl salicylaldimino])} M_2(II) in CTAB micellar solution, 453–459
Nonionic microemulsions, oil-induced structural changes in, 119–124
Nonpolar solvents, extraction and dispersion of large gold nanoparticles in, 485–489
Nucleation and growth model, balanced, for controlled precipitations, 125–138

O

Oil-induced structural changes, in nonionic microemulsions, 119–124
Oil-water-mixed surfactant systems, ultra-low interfacial tension in, 255–259
Oligodeoxyribonucleotides, activation with 2,4-phenylene diisothiocyanate and covalent grafting onto amine-functionalized latex microspheres, 473–484
Organic particles, precipitation and condensation of, 499–528
Ostwald ripening, and related phenomena, 1–12

P

Percolation, of inverted cylindrical branched micelles, 211–219
Perfluorooctanol
  phase behavior of dodecyl dimethyl amino oxide/SDS in the presence of, 111–117
  phase behavior of tetradecyldimethylaminoxide in the presence of, 431–442
Phase behavior
  of dodecyl dimethyl amino oxide/SDS in the presence of perfluorooctanol, 111–117
  of tetradecyldimethylaminoxide in the presence of perfluorooctanol and HCl, 431–442
  of Triton X-100/n- $C_8H_{17}OH$/$PEG1000_{aq}$ system, 443–451
Phase inversion, of liquid–liquid dispersions under applied electric fields, 57–69
Phenolic/furfural aerogel microspheres, direct formation in supercritical carbon dioxide, 79–87
Phenolphthalein pH-indicator, encapsulation into a sol-gel matrix, 583–590
Phenothiazine, in CTAB/n- $C_5H_{11}OH/H_2O$ microemulsion, for fluorescence quenching of anthracene, 329–335
2,4-Phenylene diisothiocyanate, for activation of oligodeoxyribonucleotides, 473–484
Photocatalytic activity, of inorganic sunscreens, 381–386
(+)-α-Pinene
  solubilization by cyclodextrin/surfactant mixed systems, 191–195
  solubilizing ability of cyclodextrins and surfactants for, 185–190
Polyacrylate dispersions, interface electrophoresis method for stability of, 291–295
Polyethylene glycol tert-octylphenyl ether solutions, concentrated, linear viscoelasticity of, 409–420
Polymerization, of cationic surfactant films in microemulsions, 597–606
Polyvinyl alcohol acetate groups, influence on adsorption and electrochemical properties of the TiO_2-polymer solution interface, 281–289
Polyvinylpyrrolidone, effect on the states of water in W/O microemulsions with betaine surfactant, 563–567
Precipitations
  controlled, a balanced nucleation and growth model for, 125–138
  of nanoscale aluminum hydroxide particles, 491–498
  of organic particles, 499–528
  seeded, in the continuous stirred tank reactor crystallizer, 373–380
Protein flocculation, structural and composition effect of anionic copolymer on, 203–209

S

Salt, influence of addition on the adsorption of a divalent cationic surfactant onto silica, 387–396
SDS. *See* Sodium dodecyl sulfate
Seeded precipitations, in the continuous stirred tank reactor crystallizer, 373–380
Short chain cosurfactant, influence of monomers on microemulsions with, 231–244
Silica
  influence of added salt on the adsorption of a divalent cationic surfactant onto, 387–396
  particles, preparation in emulsion systems, 363–371
Silicone surfactants, A-B type, solubilization enhancement effect in microemulsions, 245–253
SOD. *See* Superoxide dismutase
Sodium dodecyl sulfate (SDS)
  and dodecyl dimethyl amino oxide, phase behavior of perfluorooctanol in the presence of, 111–117
  fluorescence probe study of interaction with acrylic acid-co-ethyl methacrylate copolymers, 177–184
Sodium oleate, adsorption on kaolinite, 355–362

Sol-gel matrix, encapsulation of phenolphthalein pH-indicator into, 583–590
Solubilization, enhancement by A-B-type silicone surfactants in microemulsions, 245–253
Solubilizing ability, of cyclodextrins and surfactants for (+)-α-pinene, 185–190
Spherical particles, electrical double layer potential for, 349–354
Spontaneous emulsification, of alkyl ketene dimer, 569–581
Stability
  accelerated testing of a W/O emulsion, 23–31
  experimental design for evaluation of, 297–309
  of polyacrylate dispersion by the interface electrophoresis method, 291–295
  of W/O emulsions for food applications, polyols, high pressure, and refractive indices equalization for improvement of, 269–280
Sunscreens, inorganic, photocatalytic activity of, 381–386
Supercritical carbon dioxide, direct formation of phenolic/furfural aerogel microspheres in, 79–87
Superoxide dismutase (SOD)
  activity, effect of microenvironment created by surfactant on, 89–96
  effect of surfactant microenvironment on copper(II)-salicylate complex activity similar to, 197–202
Surfactant/alkali/acidic crude oil systems, effect of acidic components on dynamic interfacial tensions in, 41–55
Surfactants
  A-B-type silicone, solubilization enhancing effect in microemulsions, 245–253
  acid- and oxidatively-labile vinyl ether, synthesis and drug delivery applications of, 399–407
  anionic, rheological properties in the presence of Al^{3+} counterion, 421–429
  betaine, effect of polyvinylpyrrolidone on the states of water in W/O microemulsions with, 563–567
  cationic films, polymerization in microemulsions, 597–606
  and cyclodextrin mixed systems, solubilization of (+)-α-pinene by, 191–195
  derived from carboxylic acid sugars, 167–176
  divalent cationic, influence of added salt on adsorption onto silica, 387–396
  effects on hydride transfer from methylene blue leudodye to 2,5- dihydroxyl-1,4-benzoquinone in aqueous solution, 535–539
  ionic, effects of metal ions on the micellization of, 529–533
  microenvironment
    effect on SOD activity, 89–96
    effect on SOD-like activity of copper(II)-salicylate complex, 197–202
  mixed nonionic sugar based, in a ternary system with medium chain triglycerides and water, 261–268
  oil-water-mixed systems, ultra-low interfacial tensions in, 255–259
  solubilizing ability for (+)-α-pinene, 185–190

T

Ternary system, based on medium chain triglycerides/water/mixed nonionic sugar-based surfactants, 261–268
Tetradecyldimethylaminoxide, phase behavior in the presence of perfluorooctanol and HCl, 431–442
TiO_2-polymer solution interface, influence of the acetate groups from polyvinyl alcohol on adsorption and electrochemical properties of, 281–289
Triclosan-containing microcapsules, preparation and characterization by complex coacervation, 591–596
Triglycerides, medium chain, in a ternary system with mixed nonionic sugar-based surfactants, 261–268
Triton X-100/n- $C_8H_{17}OH/PEG1000_{aq}$ system, phase behavior and structural properties of, 443–451

V

Vinyl ether surfactants, acid and oxidatively labile, synthesis and drug delivery applications of, 399–407
Viscoelasticity, linear, of concentrated polyethylene glycol tert- octylphenyl ether solutions, 409–420
Viscosity, measurement for water-in-crude-oil emulsions under flowing conditions, 33–39

W

W/O. *See* Water-in-oil
W/O/W. *See* Water-in-oil-in-water
Water-in-crude-oil emulsions, viscosity measurement under flowing conditions, 33–39
Water-in-oil (W/O) emulsions
  accelerated stability testing of, 23–31
  for food applications, polyols, high pressure, and refractive indices equalization for improved stability of, 269–280
Water-in-oil-in-water (W/O/W) multiple emulsions, of calcium thioglycolate, 13–21